

# NLO corrections to $\Delta F = 2$ effective Hamiltonians and SUSY contribution to FCNC

Ignazio Scimemi<sup>a</sup>

<sup>a</sup>Dip. di Fisica, Università degli Studi di Roma “La Sapienza”, P.le A. Moro 2, 00185 Roma, Italy.

The most general QCD-NLO Anomalous-Dimension Matrix of all four-fermion dimension six  $\Delta F = 2$  operators is presented. Two applications of this Anomalous-Dimension Matrix to the study of SUSY contribution to  $K - \bar{K}$  mixing are also discussed.

## 1. Introduction

Theoretical predictions of several measurable quantities, which are relevant in  $K$ -,  $D$ - and  $B$ -meson phenomenology, depend on the matrix elements of some  $\Delta F = 2$  four-fermion operators. Examples are given by FCNC effects in SUSY extensions of the Standard Model [1]-[2], by the  $B_s^0 - \bar{B}_s^0$  width difference [3] and by the  $O(1/m_b^3)$  corrections in inclusive b-hadron decay rates (which actually depend on the matrix elements of several four-fermion  $\Delta F = 0$  operators [4]). In all these cases, the relevant operators have the form

$$Q = C^{\alpha\beta\rho\sigma} (\bar{b}_\alpha \Gamma q_\beta) (\bar{b}_\rho \Gamma q_\sigma), \quad (1)$$

where  $\Gamma$  is a generic Dirac matrix acting on (implicit) spinor indices;  $\alpha - \sigma$  are colour indices and  $C^{\alpha\beta\rho\sigma}$  is either  $\delta^{\alpha\beta} \delta^{\rho\sigma}$  or  $\delta^{\alpha\sigma} \delta^{\rho\beta}$ . All the operators discussed in this paper appear in some “effective” theory, obtained by using the Operator Product Expansion (OPE). As a consequence, in all cases, three steps are necessary for obtaining physical amplitudes from their matrix elements:

- i) matching of the original theory to the effective one at some large energy scale;
- ii) renormalization-group evolution from the large energy scale to a low scale suitable for the calculation of the hadronic matrix elements (typically 1–5 GeV);
- iii) non-perturbative calculation of the hadronic matrix elements.

In the following a calculation of the two-loop Anomalous Dimension Matrix (ADM) relevant for  $\Delta F = 2$  transition amplitudes and its application to FCNC in SUSY extension of the Stan-

dard Model (SM) is presented. I refer to the original papers for all details of the calculations, *i.e.*, respectively to ref. [5] for the calculation of the ADM and to ref. [6,7] for its applications to FCNC effects in Supersymmetry.

The paper is organized as follows. In sec. 2 I introduce the operator basis, the Wilson coefficients and the Renormalization Group Equations (RGE). In sec. 3 constraints on SUSY parameters coming from FCNC in the  $K - \bar{K}$  mixing are derived.

## 2. Effective Hamiltonian formalism and ADM

The three steps (i-ii-iii, sec.1) needed to use OPE are treated in the following.

In all cases of interest, the matrix elements of the effective Hamiltonian can be written as

$$\langle F | \mathcal{H}_{eff} | I \rangle = \sum_i \langle F | Q_i(\mu) | I \rangle C_i(\mu), \quad (2)$$

where the  $Q_i(\mu)$ s are the relevant operators renormalized at the scale  $\mu$  and the  $C_i(\mu)$ s are the corresponding Wilson coefficients.

The set of operators relevant for a description of FCNC effects in SUSY models is:

$$\begin{aligned} Q_1 &= \bar{d}^\alpha \gamma_\mu (1 - \gamma_5) s^\alpha \bar{d}^\beta \gamma_\mu (1 - \gamma_5) s^\beta, \\ Q_2 &= \bar{d}^\alpha (1 - \gamma_5) s^\alpha \bar{d}^\beta (1 - \gamma_5) s^\beta, \\ Q_3 &= \bar{d}^\alpha (1 - \gamma_5) s^\beta \bar{d}^\beta (1 - \gamma_5) s^\alpha, \\ Q_4 &= \bar{d}^\alpha (1 - \gamma_5) s^\alpha \bar{d}^\beta (1 + \gamma_5) s^\beta, \\ Q_5 &= \bar{d}^\alpha (1 - \gamma_5) s^\beta \bar{d}^\beta (1 + \gamma_5) s^\alpha, \end{aligned} \quad (3)$$

together with operators  $\tilde{Q}_{1,2,3}$  which can be ob-

tained from  $Q_{1,2,3}$  by the exchange  $(1 - \gamma_5) \leftrightarrow (1 + \gamma_5)$ .

Let us now represent the operators as row vectors  $\vec{Q}$  and the coefficients,  $\vec{C}(\mu)$ , as column ones. The vectors  $\vec{C}(\mu)$  are expressed in terms of their counter-part, computed at a large scale  $M$ , through the renormalization-group evolution matrix  $\hat{W}[\mu, M]$

$$\vec{C}(\mu) = \hat{W}[\mu, M] \vec{C}(M). \quad (4)$$

The initial conditions for the evolution equations,  $\vec{C}(M)$ , are obtained by matching the full theory, which includes propagating heavy-vector bosons ( $W$  and  $Z^0$ ), the top quark, SUSY particles, etc., to the effective theory where the  $W$ ,  $Z^0$ , the top quark and all the heavy particles have been removed. In general,  $\vec{C}(M)$  depend on the definition of the operators in a given renormalization scheme.

The coefficients  $\vec{C}(\mu)$  obey the renormalization-group equations:

$$\left[ -\frac{\partial}{\partial t} + \beta(\alpha_s) \frac{\partial}{\partial \alpha_s} + \beta_\lambda(\alpha_s) \lambda \frac{\partial}{\partial \lambda} - \frac{\hat{\gamma}^T(\alpha_s)}{2} \right] \vec{C}(t, \alpha_s(t), \lambda(t)) = 0, \quad (5)$$

where  $t = \ln(M^2/\mu^2)$ . The term proportional to  $\beta_\lambda$ , the  $\beta$ -function of the gauge parameter  $\lambda(t)$  (for covariant gauges), takes into account the gauge dependence of the Wilson coefficients in gauge-dependent renormalization schemes, such as the RI scheme [10]. This term is absent in standard  $\overline{\text{MS}}$  schemes, independently of the regularization which is adopted (NDR, HV or DRED for example).

If no quark threshold between the scales  $M$  and  $\mu$  is crossed, eq. (5) can be easily solved (the formulae for the general case can be found in refs. [11,12]). At the next-to-leading order, one can write

$$\hat{W}[\mu, M] = \hat{M}[\mu] \hat{U}[\mu, M] \hat{M}^{-1}[M], \quad (6)$$

where  $\hat{U}$  is the leading-order evolution matrix

$$\hat{U}[\mu, M] = \left[ \frac{\alpha_s(M)}{\alpha_s(\mu)} \right]^{\hat{\gamma}^{(0)T}/2\beta_0}, \quad (7)$$

and the NLO matrix is given by

$$\hat{M}[\mu] = \hat{1} + \frac{\alpha_s(\mu)}{4\pi} \hat{J}[\lambda(\mu)]. \quad (8)$$

By substituting the expression of the  $\vec{C}(\mu)$  given in eq. (4) in the renormalization-group equations (5), and using  $\hat{W}[\mu, M]$  written as in eqs. (6)–(8), one finds that the matrix  $\hat{J}$  satisfies the equation

$$\hat{J} + \frac{\beta_\lambda^0}{\beta_0} \lambda \frac{\partial \hat{J}}{\partial \lambda} - \left[ \hat{J}, \frac{\hat{\gamma}^{(0)T}}{2\beta_0} \right] = \frac{\beta_1}{2\beta_0^2} \hat{\gamma}^{(0)T} - \frac{\hat{\gamma}^{(1)T}}{2\beta_0}. \quad (9)$$

In eqs. (7) and (9),  $\beta_0$ ,  $\beta_1$  and  $\beta_\lambda^0$  are the first coefficients of the  $\beta$ -functions of  $\alpha_s$  and of  $\lambda$ , respectively;  $\hat{\gamma}^{(0)}$  and  $\hat{\gamma}^{(1)}$  are the LO and NLO anomalous dimension matrices.  $\hat{U}$  is determined by the LO anomalous dimension matrix  $\hat{\gamma}^{(0)}$  and is therefore regularization and renormalization-scheme independent; the two-loop anomalous dimension matrix  $\hat{\gamma}^{(1)}$ , and consequently  $\hat{J}$  and  $\hat{W}[\mu, M]$ , are, instead, renormalization-scheme dependent. The precise definition and a complete discussion of the scheme dependence of eq. (9) can be found in ref. [5] and in references therein.

Eventually, the B-parameters which parametrize the hadronic matrix elements of eq. (3) have been evaluated non-perturbatively on the lattice [13,14].

In what follows I have used the estimate of ref. [13].

### 3. FCNC in SUSY

The considered operator basis is the one relevant for the analysis of FCNC in SUSY in the so-called mass insertion approximation [8]. In this approach for fermions and sfermions states all the couplings of these particles to neutral gauginos are flavor diagonal and FCNC effects are shown by the non-diagonality of sfermion propagators. The pattern of flavor change, for the K-system, is given by the ratio

$$(\delta_{ij}^d)_{AB} = \frac{(m_{ij}^{\tilde{d}})_{AB}^2}{M_{sq}^2}, \quad (10)$$

where  $(m_{ij}^{\tilde{d}})_{AB}^2$  are the off-diagonal elements of the  $\tilde{d}$  mass squared matrix that mixes fla-

for  $i, j$  for both left- and right-handed scalars ( $A, B = \text{Left, Right}$ ), and  $M_{sq}$  is an average squark mass, see *e.g.* [1]. The sfermion propagators are expanded as a series in terms of the  $\delta$ 's and the contribution of the first term of this expansion is considered. Assuming that all s-particles are heavier than fermions one produces effective four fermions interactions and OPE can be used.

The Wilson coefficients at the matching scale  $M_{sq}$  can be found, *e.g.*, in ref. [1,6,7,9] and I refer to these articles for their complete expression.

One can provide a set of constraints on SUSY variables coming from the  $K_L - K_S$  mass difference,  $\Delta M_K$  and the CP violating parameter  $\epsilon_K$  defined as

$$\begin{aligned}\Delta M_K &= 2\text{Re}\langle K^0 | H_{\text{eff}} | \bar{K}^0 \rangle, \\ \epsilon_K &= \frac{1}{\sqrt{2}\Delta M_K} \text{Im}\langle K^0 | H_{\text{eff}} | \bar{K}^0 \rangle.\end{aligned}\quad (11)$$

The parameter space is composed of two real and four complex entries, that is  $M_{sq}$ ,  $m_{\tilde{g}}$  and  $(\delta_{12}^d)_{LL}$ ,  $(\delta_{12}^d)_{LR}$ ,  $(\delta_{12}^d)_{RL}$ ,  $(\delta_{12}^d)_{RR}$ .

Neglecting interference among different SUSY contribution one gives upper bound on the  $\delta$ 's, giving specific values to  $M_{sq}$  and  $m_{\tilde{g}}$ . I consider the conditions  $M_{sq} \sim m_{\tilde{g}}$  in subsec. 3.1 and the case in which the scalars of the first-two generations are heavier than the rest of the supersymmetric spectrum in subsec. 3.2. Constraints on individual  $\delta$ 's are provided. Indeed, it is meaningful to study the interference of cancellation effects only in specific models.

The physical condition used to get the bounds on the  $\delta$ 's is that the SUSY contribution (proportional to each single  $\delta$ ) plus the SM contribution to  $\Delta M_K$  and  $\epsilon_K$  do not exceed the experimental value of these quantities.

### 3.1. $M_{sq} \sim m_{\tilde{g}}$

For Left-Right mass insertions, I consider two possible (extreme) cases:  $|(\delta_{12}^d)_{LR}| \gg |(\delta_{12}^d)_{RL}|$  and  $|(\delta_{12}^d)_{LR}| = |(\delta_{12}^d)_{RL}|$ . In the second case the contributions proportional to  $(\delta_{12}^d)_{LR}^2$ ,  $(\delta_{12}^d)_{RL}^2$  and  $(\delta_{12}^d)_{LR}(\delta_{12}^d)_{LR}$  are combined. This approach improves the one of refs. [9,15], where contributions proportional to  $(\delta_{12}^d)_{LR}(\delta_{12}^d)_{LR}$  were considered independently from the ones proportional to

$(\delta_{12}^d)_{LR}^2$  and  $(\delta_{12}^d)_{RL}^2$ .

$(\delta_{12}^d)_{LL}(\delta_{12}^d)_{RR}$  contribution can be treated independently from the  $(\delta_{12}^d)_{LL}^2$  and  $(\delta_{12}^d)_{RR}^2$  ones, since the latter ones do not generate Left-Right operators and they are suppressed with respect to the first contribution.

Results for the bounds of the  $\delta$ 's for  $M_{sq} \sim 200$  GeV and various values of the ratio  $x = m_{\tilde{g}}^2/M_{sq}^2$  derived for real parts of the  $\delta$ 's are shown in tab. 1. Let us compare these results with the previous analyses of refs. [1,9]. In both papers the matrix elements were computed in the Vacuum Insertion Approximation (VIA) and with no evolution [1] or leading order evolution [9] of the coefficient functions. In this analysis lattice B-parameters and NLO evolution of the coefficients have been calculated. This allows a consistent matching of the renormalization-scheme and  $\mu$  dependence of the QCD-corrected  $\mathcal{H}_{eff}^{\Delta S=2}$ .

### 3.2. Heavy first-two generation scalars

Let us assume now that the scalars of the first two generations (the typical mass of which is here again denoted by  $M_{sq}$ ) is much heavier than the rest of the SUSY spectrum (which has a typical mass of  $m_{\tilde{g}}$ ). In this case the heavy squarks have to be integrated out as a first, at a scale  $M_{sq}$ . Then, the coefficients must be evolved to the scale  $m_{\tilde{g}}$  where all the rest of the supersymmetric particles are also integrated out.

In table 2 results are shown for the real parts of the  $\delta$ 's with  $m_{\tilde{g}} \sim 250$  GeV and several values of  $M_{sq}$ .

The constraints that come from the four possible insertions of the  $\delta$ 's are presented: in the first and second rows only terms proportional respectively to  $(\delta_{12}^d)_{LL}$  and  $(\delta_{12}^d)_{LR}$  are considered; in the last two rows the contribution of operators with opposite chirality,  $RR$  and  $RL$ , is also evaluated by assuming  $(\delta_{12}^d)_{LL} = (\delta_{12}^d)_{RR}$  and  $(\delta_{12}^d)_{LR} = (\delta_{12}^d)_{RL}$ .

The theoretical improvements of this results with respect to ref.[9] have already been discussed in subsec. 3.1. The combination of B-parameters and NLO-QCD corrections change the LO-VIA results by about 25 – 35%. As expected [9], the tightest constraints are for the cases  $(\delta_{12}^d)_{LL} = (\delta_{12}^d)_{RR}$  and  $(\delta_{12}^d)_{LR} = (\delta_{12}^d)_{RL}$ . In these cases

|     | No-QCD,<br>VIA<br>$\times 10^{-3}$                                                                   | LO,<br>VIA<br>$\times 10^{-3}$ | LO,<br>$B_i$<br>$\times 10^{-3}$ | NLO,<br>$B_i$<br>$\times 10^{-3}$ |
|-----|------------------------------------------------------------------------------------------------------|--------------------------------|----------------------------------|-----------------------------------|
| $x$ | $\sqrt{ \text{Re}(\delta_{12}^d)_{LL}^2 }$                                                           |                                |                                  |                                   |
| 0.3 | 5.0                                                                                                  | 5.7                            | 7.7                              | 7.7                               |
| 1.0 | 11                                                                                                   | 12                             | 16                               | 16                                |
| 4.0 | 25                                                                                                   | 29                             | 39                               | 39                                |
| $x$ | $\sqrt{ \text{Re}(\delta_{12}^d)_{LR}^2 } \quad ( (\delta_{12}^d)_{LR}  \gg  (\delta_{12}^d)_{RL} )$ |                                |                                  |                                   |
| 0.3 | 1.1                                                                                                  | 0.84                           | 1.1                              | 0.96                              |
| 1.0 | 1.2                                                                                                  | 0.93                           | 1.2                              | 1.1                               |
| 4.0 | 1.8                                                                                                  | 1.3                            | 1.6                              | 1.5                               |
| $x$ | $\sqrt{ \text{Re}(\delta_{12}^d)_{LR}^2 } \quad ((\delta_{12}^d)_{LR} = (\delta_{12}^d)_{RL})$       |                                |                                  |                                   |
| 0.3 | 2.0                                                                                                  | 1.4                            | 0.89                             | 0.67                              |
| 1.0 | 1.1                                                                                                  | 0.97                           | 1.8                              | 3.0                               |
| 4.0 | 1.3                                                                                                  | 1.0                            | 1.4                              | 1.3                               |
| $x$ | $\sqrt{ \text{Re}(\delta_{12}^d)_{LL}(\delta_{12}^d)_{RR} }$                                         |                                |                                  |                                   |
| 0.3 | 0.64                                                                                                 | 0.39                           | 0.40                             | 0.33                              |
| 1.0 | 0.71                                                                                                 | 0.44                           | 0.45                             | 0.37                              |
| 4.0 | 1.0                                                                                                  | 0.61                           | 0.62                             | 0.52                              |

Table 1

Limits on  $\text{Re}(\delta_{ij})_{AB}(\delta_{ij})_{CD}$ , with  $A, B, C, D = (L, R)$ , for an average squark mass  $M_{sq} = 200$  GeV and for different values of  $x = m_g^2/M_{sq}^2$ .

the coefficients proportional to  $(\delta_{12}^d)_{LL}(\delta_{12}^d)_{RR}$ ,  $(\delta_{12}^d)_{LR}(\delta_{12}^d)_{RL}$  dominate the others.

#### 4. Acknowledgements

I wish to thank all the authors of refs. [5–7] with whom the results presented in this work have been obtained.

Particular thanks to Stephan Narison for the beautiful conference he has organized.

#### REFERENCES

1. E. Gabrielli, A. Masiero and L. Silvestrini, Phys. Lett. B374 (1996) 80; F. Gabbiani, E. Gabrielli, A. Masiero and L. Silvestrini, Nuc. Phys. B477 (1996) 321 and refs. therein.
2. J.A. Bagger, K.T. Matchev and R.J. Zhang, preprint JHU-TIPAC-97011, hep-ph/9707225 and refs. therein.
3. M. Beneke, G. Buchalla and I. Dunietz, Phys. Rev. D54 (1996) 4419 and refs. therein.

| $M_{sq}$<br>[TeV] | No-QCD,<br>VIA<br>$\times 10^{-2}$                                           | LO,<br>VIA<br>$\times 10^{-2}$ | LO,<br>$B_i$<br>$\times 10^{-2}$ | NLO,<br>$B_i$<br>$\times 10^{-2}$ |
|-------------------|------------------------------------------------------------------------------|--------------------------------|----------------------------------|-----------------------------------|
|                   | $\sqrt{ \text{Re}(\delta_{12}^d)_{LL}^2 }$                                   |                                |                                  |                                   |
| 2                 | 3.1                                                                          | 3.6                            | 4.9                              | 4.9                               |
| 5                 | 7.5                                                                          | 8.8                            | 0.12                             | 0.12                              |
| 10                | 15                                                                           | 18                             | 25                               | 24                                |
|                   | $\sqrt{ \text{Re}(\delta_{12}^d)_{LR}^2 }$                                   |                                |                                  |                                   |
| 2                 | 2.1                                                                          | 1.4                            | 1.8                              | 1.6                               |
| 5                 | 9.8                                                                          | 6.5                            | 8.2                              | 7.2                               |
| 10                | 34                                                                           | 22                             | 28                               | 25                                |
|                   | $\sqrt{ \text{Re}(\delta_{12}^d)_{LR} = \text{Re}(\delta_{12}^d)_{RL} }$     |                                |                                  |                                   |
| 2                 | 0.66                                                                         | 0.35                           | 0.35                             | 0.28                              |
| 5                 | 1.5                                                                          | 0.77                           | 0.79                             | 0.64                              |
| 10                | 3.0                                                                          | 1.5                            | 1.5                              | 1.2                               |
|                   | $\sqrt{ \text{Re}(\delta_{12}^d)_{LL}^2 = \text{Re}(\delta_{12}^d)_{RR}^2 }$ |                                |                                  |                                   |
| 2                 | 1.1                                                                          | 0.52                           | 0.51                             | 0.41                              |
| 5                 | 4.1                                                                          | 1.6                            | 1.6                              | 1.3                               |
| 10                | 10                                                                           | 3.6                            | 3.4                              | 2.7                               |

Table 2

Limits on  $\text{Re}(\delta_{12}^d)_{AB}$  from  $\Delta M_K$  with gaugino masses of 250 GeV.

4. M. Neubert and C.T. Sachrajda, Nucl. Phys. B483 (1997) 339 and refs. therein.
5. M. Ciuchini et al. Nuc. Phys. B523 (1998) 501.
6. M. Ciuchini et al. hep-ph/9808328.
7. R. Contino and I. Scimemi, preprint ROME1-1216/98.
8. L. J. Hall, V. A. Kostolecki and S. Raby, Nucl. Phys. B267 (1986) 415.
9. J. A. Bagger, K. T. Matchev and R. J. Zhang, Phys. Lett. B412 (1997) 77.
10. M. Ciuchini *et al.*, Z. Phys. C68 (1995) 239.
11. M. Ciuchini, E. Franco, G. Martinelli and L. Reina, Nucl. Phys. B415 (1994) 403.
12. A.J. Buras, M. Jamin, M.E. Lautenbacher and P.H. Weisz, Nucl. Phys. B370 (1992) 69; Addendum, Nucl. Phys. B375 (1992) 501.
13. C. R. Allton et al. hep-lat/9806016
14. R. Gupta, T. Bhattacharya and S. Sharpe, Phys. Rev. D55 (1997) 4036.
15. J.S. Hagelin, S. Kelley and T. Tanaka, Nucl. Phys. B322 (1989) 55.